

LAURENCE E. STAWICK, M.D., A.G.A.F. MARK S. DeVORE, M.D., F.A.C.P. JULIA S. GREER, M.D. KHA H. NGO, D.O. SERGE A. SORSER, M.D. Phone (248) 662-4110 Fax (248) 662-4120 Phone (248) 569-1770 Fax (248) 443-2439

Miralax/Gatorade Bowel Preparation

Importance of DRINKING LIQUIDS during the bowel preparation process

During bowel preparation you will lose significant amounts of fluid. THIS IS NORMAL. It is very important that you replace this fluid to prevent dehydration. Drink large amounts of clear liquids. Drinking large amounts of clear liquids also helps ensure that your bowel will be clean for the examination. A completely clean colon may help avoid the need for a repeat exam.

Read carefully. Follow the steps listed below

Purchase both at the Drugstore.

(Over the counter) Miralax: 238 grams. 4 Dulcolax 5mg Tablets 64oz Gatorade, Propel, water or other clear liquid, Diabetic patients- use crystal light

DAY BEFORE EXAM 8:00 a.m. Do not eat until after your procedure, Clear liquids only (see list)⇒ Keep Hydrated all day. Avoid dairy products







ALL DAY LONGCLEAR LIQUIDS DIET LIST
(DO NOT DRINK ANYTHING COLORED RED OR PURPLE)Water, tea or coffee (no milk or non-dairy
creamer) sweeteners are o.k.Soft drinks (7-up, cola, ginger ale, orange, Sprite,
etc.) Gatorade, Propel, Kool-Aid, lemonade.Strained fruit juices; without pulp.
(apple, white cranberry, orange, white grape, etc.)Low sodium chicken or beef bouillon/broth
Hard candies, Jell-O, Popsicles
NO sherbets or fruit bars

Your body loses significant amounts of fluid during bowel preparation. In order to prevent dehydration, it is important to supplement that fluid loss with clear liquids. Make a conscious effort to drink as much as you can before, during, and after the preparation.

Evening before procedure, Split Dose (2-Day) Regimen
2:00 p.m. take 2 Dulcolax 5mg laxative tablets Mix the 238g bottle of Miralax in 64 ounces of cold clear liquid of your choice. (Mix entire bottle) POP and juice not recommended

4:00 p.m. First dose: (You may vary times by 2 hours - between 4:00 to 6:00 pm) Start drinking the Miralax mix. Drink 4 glasses. Drink 1 (8 oz. glasses) every 30 minutes.

DAY OF EXAM

5 hours prior to your procedure

Take 2 Dulcolax 5mg laxative tablets Start drinking the Miralax mix. Drink 4 glasses. Drink 1 (8 oz. glasses) every 30 minutes until gone.

Do not drink anything 3 hours prior to your procedure on the morning of !!!

For coupons online see website: https://www.miralax.com/coupons-and-offers